

### FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

Name: D.O.B.:	PLACE PICTURE HERE					
Weight:Ibs. Asthma: [ ] Yes (higher risk for a severe reaction) [ ] No						
NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.						
Extremely reactive to the following foods:						
THEREFORE: [ ] If checked, give epinephrine immediately for ANY symptoms if the allergen was likely eaten.						
[ ] If checked, give epinephrine immediately if the allergen was definitely eaten, even if no symptoms are noted.						

#### FOR ANY OF THE FOLLOWING:

# **SEVERE** SYMPTOMS



Short of breath, wheezing, repetitive cough



Pale, blue, faint, weak pulse, dizzy



Tight, hoarse, trouble breathing/ swallowing



Significant swelling of the tongue and/or lips

ORA

from different

body areas.



SKIN

Many hives over body, widespread redness



Repetitive vomiting, severe diarrhea

Ţ



Feeling something bad is about to happen, anxiety, confusion





OTHER



1

#### 1. INJECT EPINEPHRINE IMMEDIATELY.

2. Call 911. Tell them the child is having anaphylaxis and may need epinephrine when they arrive.

Ð.

- Consider giving additional medications following epinephrine:
  - Antihistamine
  - Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport them to ER even if symptoms resolve. Person should remain in ER for at least 4 hours because symptoms may return.

# **MILD** SYMPTOMS









Itchy/runny Itchy mouth nose, sneezing

A few hives. mild itch

Mild nausea/ discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

#### FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA. FOLLOW THE DIRECTIONS BELOW:

- 1. Antihistamines may be given, if ordered by a healthcare provider.
- 2. Stay with the person; alert emergency contacts.
- 3. Watch closely for changes. If symptoms worsen, give epinephrine.

6.4	1 <b>^</b> A T	IAAL	らんしつ	SES
IVI	ILAI	IUN	อเมน	JOES

Epinephrine Brand:							
Epinephrine Dose: [ ] 0.15 mg IM [ ] 0.3 mg IM							
Antihistamine Brand or Generic:							
Antihistamine Dose:							
Other (e.g., inhaler-bronchodilator if wheezing):							

# ST. LUKE'S SCHOOL PEANUT/NUT & FOOD ALLERGY INDIVIDUAL HEALTH CARE PLAN (IHCP)

Name:		_ DOB:	Allergy to	o:			
Level of Allergy: (Circle	e) Inhalation Tactile	Ingestion	Unknown	Age of Onset:			
Describe symptom(s)	of allergic reaction(s):						
History of Anaphylaxis	:: (Circle one) Yes No T	reatment: EpiP	en Benadryl	Other: (please sp	ecify)		
Other health condition	ns/medications (e.g., Asth	ma):			n		
	ne at school: Health Office	e (unlocked cab	inet during reg	ular school hours	)		
Preventive Measu Signs shall be posted a	res Idvising there is a student	t with allergies	to peanuts/nu	ts.	Parent/Guardian	<u>Initials</u>	
Student needs a designated peanut/nut free table or desk in the classroom and lunch table. Initial - Yes No  *Before participation, the parent/guardian will inform the food service of their child's food allergy*  Student will only eat foods sent in from his/her own home for class parties and snacks. Initial - Yes No  Parent/guardian will provide a safe snack for class parties to be kept at school. Initial - Yes No  Parent/guardian will inform bus company personnel of their child's allergy medication requirements, & emergency contact information.  Parent/Guardian Initials  Medications that are kept in the nurse's office are available during school hours only. For any before and after school activities (including C.A.M.P.), it is the parent/guardian responsibility to inform activity coordinator of their child's allergy, treatments & provide emergency medications.  Parent/Guardian Initials  School personnel who may be involved in the care of a student who has been diagnosed with an allergy will be informed of the EHCP and IHCP.  All parents/guardians in the peanut/nut allergy student's classroom will be notified in writing by the school administrator of the peanut/nut free snack policy.  Student will be reminded not to share or trade food.  The student will be accompanied to the health office in the event of an allergic reaction  Trained school personnel will carry the prescribed emergency medication(s), and a copy of the EHCP on the field trip and accompany student at all times if the parent is not present. NOTE: Student may self-carry with doctor orders.  Other Provisions Required:							
Parent/Guardian:	NTACTS (please print)			Phone: ()	<del></del>	LL 911	
	onship:						
Physician/HCP Autho	rization Signature			Date	<del></del>		
Parent/Guardian Aut	horization Signature			Date			
Certified School Nurs	e Teacher Authorization S	ignature		Date			